

## Tutorials 6 – SDE (2) : Anderson localization in 1D

### 1 Disordered Schrödinger equation & Anderson localization

We consider the one-dimensional Schrödinger equation with a disordered potential

$$-\psi''(x) + V(x)\psi(x) = E\psi(x). \quad (1)$$

We choose the simplest model of disorder and assume that  $V(x)$  is a Gaussian white noise (in space)

$$\langle V(x) \rangle = 0 \quad \text{and} \quad \langle V(x)V(x') \rangle = \sigma \delta(x - x'), \quad (2)$$

where  $\sigma$  measures the disorder strength. The aim of the exercise is to study the statistical properties of the wave function  $\psi(x)$ . In the following we study the Cauchy (initial value) problem, i.e. the solution obtained for given initial conditions  $\psi(0)$  and  $\psi'(0)$  (and not the Sturm-Liouville (spectral) problem defined by boundary conditions  $\psi(0)$  and  $\psi(L)$ ).

- 1/ We have chosen units such that  $\hbar^2/(2m) = 1$ , hence all dimensions can be expressed in terms of length. What is the dimension of an energy ? And the disorder strength  $\sigma$  ?
- 2/ **Prüfer variables.**— We consider the solution of energy  $E = k^2 > 0$  such that  $\psi(0) = 0$  and  $\psi'(0) = k$ . Write two first order linear differential equations for  $\psi$  and  $\psi'$ . Performing the change of variables  $\psi(x) = \rho(x) \sin \theta(x)$  and  $\psi'(x) = k \rho(x) \cos \theta(x)$ , show that  $\rho$  and  $\theta$  obey

$$\frac{d\theta(x)}{dx} = k - \frac{V(x)}{k} \sin^2 \theta(x) \quad (\text{Stratonovich}) \quad (3)$$

$$\frac{d \ln \rho(x)}{dx} = \frac{V(x)}{2k} \sin 2\theta(x) \quad (\text{Stratonovich}) \quad (4)$$

What are the initial conditions  $\rho(0)$  and  $\theta(0)$  ?

- 3/ Introduce  $\xi(x) = \ln \rho(x)$  and give the two Itô SDE for  $\theta(x)$  and  $\xi(x)$  : you can write  $V(x)dx = \sqrt{\sigma} dW(x)$  where  $W(x)$  is a Wiener process.
- 4/ **Localization length.**— The localization length  $\xi_{\text{loc}}$  is the characteristic length controlling the exponential growth (or decay) of the wave function. We define it as

$$1/\xi_{\text{loc}} \stackrel{\text{def}}{=} \gamma = \lim_{x \rightarrow \infty} \frac{\ln \rho(x)}{x} = \frac{d}{dx} \langle \xi(x) \rangle \quad (5)$$

where  $\gamma$  is the “Lyapunov exponent”. Which SDE (Stratonovich or Itô) is more convenient in order to get  $\gamma$  ? Assuming the phase uniformly distributed in the limit Energy  $\gg$  disorder, deduce the energy dependence of the localization length in this limit.

- 5/ **Localization of electromagnetic waves.**— Consider now the Helmholtz equation for an electromagnetic wave in a random medium

$$E''(x) + k^2 \left( 1 + \frac{\delta\epsilon(x)}{\bar{\epsilon}} \right) E(x) = 0 \quad (6)$$

where  $\delta\epsilon(x)$  represents fluctuations of the dielectric constant. Transpose the results obtained for the Schrödinger equation to this case.

## 2 Distribution of the transmission probability for disordered wave equations

We consider the transmission of a wave through a one-dimensional disordered medium and derive the distribution of the transmission probability. The nature of the wave plays no role here.




Figure 1: *Transmission of a wave through a disordered region.*

- 1/ **Preliminary.**— Given the Itô SDE  $dx(t) = a(x) dt + b(x) dW(t)$ , what is  $\langle dx(t) \rangle$  and what is  $\langle dx(t)^2 \rangle$  ?
- 2/ **Composition rule.**— Scattering of a wave through a certain region is characterized by two sets of left and right reflection/transmission amplitudes  $(r, t)$  and  $(r', t')$ , respectively (if the wave is incoming from the right, the reflected amplitude is  $r'$  and the transmission amplitude  $t'$ ). Show that the composition rule for transmission *amplitudes* of two regions is :

$$t_{2\oplus 1} = t_2 t_1 + t_2 (r'_1 r_2) t_1 + \dots = \frac{t_2 t_1}{1 - r'_1 r_2} \quad (7)$$

- 3/ **Evolution of the transmission.**— We denote  $\tau(x) = |t_1|^2 = |t'_1|^2$  the transmission *probability* of region 1 (corresponding to the interval  $[0, x]$ ). We consider a small slice of disordered medium in  $[x, x+\delta x]$ , described by reflection and transmission amplitudes  $(r_2, t_2)$  and  $(r'_2, t'_2)$ . We introduce the reflection probability  $\rho = |r_2|^2 \ll 1$ . Eq. (7) gives

$$\begin{aligned} \tau(x + \delta x) &= \frac{\tau(x)(1 - \rho)}{|1 + e^{i\phi} \sqrt{1 - \tau(x)} \sqrt{\rho}|^2} \\ &\simeq \tau(x) - 2 \cos(\phi) \tau \sqrt{1 - \tau} \sqrt{\rho} + [-\tau(2 - \tau) + 4\tau(1 - \tau) \cos^2(\phi)] \rho + \mathcal{O}(\rho^{3/2}) \end{aligned} \quad (8)$$

Assumptions :

- $\langle \rho \rangle \simeq \delta x / \ell$ , where  $\ell$  is the scattering length (an effective parameter characterising the strength of the disorder).
- The phase  $\phi$  is independent of  $\tau(x)$  and  $\rho$  and uniformly distributed (of course these assumption are not exact).

Denoting  $\delta\tau(x) = \tau(x + \delta x) - \tau(x)$ , express  $\langle \delta\tau \rangle$  and  $\langle \delta\tau^2 \rangle$  in terms of averages of functions of  $\tau$ . Deduce that the transmission obeys

$$d\tau(x) = -\tau^2 \frac{dx}{\ell} + \sqrt{\frac{2}{\ell}} \tau^2 (1 - \tau) dW(x) \quad (\text{Itô}) \quad (9)$$

- 4/ **Lyapunov exponent.**— Using Itô calculus, give  $d \ln \tau(x)$ . Deduce the relation between the effective parameter  $\ell$  and the Lyapunov exponent  $\gamma$  introduced in the first exercice.

5/ **Distribution of the transmission probability.**— We parametrise the transmission probability as  $\tau(x) = 1/\cosh^2 u(x)$ . Show that the noise becomes additive

$$du(x) = \frac{\gamma}{\tanh 2u} dx - \sqrt{\gamma} dW(x) \quad (10)$$

Considering the limit of large  $x$ , simplify the SDE and deduce the distribution of  $u(x)$ . Deduce the corresponding distribution of  $\ln \tau(L)$ , where  $\tau(L)$  is the transmission probability of a disordered region of length  $L$ . Compare the mean value and the variance.

6/ The above calculation is adapted from the famous article [2]. The *ad hoc* hypothesis made above is equivalent to the **Single Parameter Scaling** hypothesis of the gang of four [1]. In the article [3], we have compared (analytically and numerically) the two first cumulants of the log of the wave function,  $\gamma_1 = \lim_{x \rightarrow \infty} \frac{1}{x} \langle \ln |\psi(x)| \rangle$  and  $\gamma_2 = \lim_{x \rightarrow \infty} \frac{1}{x} \text{Var}(\ln |\psi(x)|)$  for the disordered model introduced in the first exercise. The result is plotted on the Figure 2. Discuss the relation with the previous results.




Figure 2: Left : The two first cumulants of  $\ln |\psi(x)|$  for  $\sigma = 1$  ( $\gamma_1$  in red and  $\gamma_2$  in blue). Right : Ratio as a function of the energy. From [3].

## Appendix :

**Beta function.**— The Beta function is defined as  $B(\mu, \nu) = \frac{\Gamma(\mu)\Gamma(\nu)}{\Gamma(\mu+\nu)}$  Useful integrals :

$$B(\mu, \nu) = \int_0^1 dt t^{\mu-1} (1-t)^{\nu-1} = 2 \int_0^{\pi/2} d\theta \sin^{2\mu-1} \theta \cos^{2\nu-1} \theta. \quad (11)$$

**Itô-Stratonovich.**— The Itô SDE  $dx_i = a_i(\vec{x}) dt + b_{ij}(\vec{x}) dW_j(t)$  and the stratonovich SDE  $dx_i = \alpha_i(\vec{x}) dt + b_{ij}(\vec{x}) dW_j(t)$  describe the *same* process if  $\alpha_i = a_i - \frac{1}{2} b_{jk} \partial_j b_{ik}$ .

## References

- [1] E. Abrahams, P. W. Anderson, D. C. Licciardello and T. V. Ramakrishnan, Scaling theory of localization: absence of quantum diffusion in two dimensions, Phys. Rev. Lett. **42**(10), 673 (1979).
- [2] P. W. Anderson, D. J. Thouless, E. Abrahams and D. S. Fisher, New method for a scaling theory of localization, Phys. Rev. B **22**(8), 3519–3526 (1980).
- [3] K. Ramola and C. Texier, Fluctuations of random matrix products and 1D Dirac equation with random mass, J. Stat. Phys. **157**(3), 497–514 (2014).