

exam 2018-2019

Renormalization à la Migdal-Kadanoff - short correction

A- The 1d case

- 1) We have $\exp(KS_iS_j) = \cosh K + S_iS_j \sinh K = (\cosh K)(1 + S_iS_j \tanh K)$. This stems from the fact that $S_iS_j = \pm 1$. If this is not seen immediately, one way to proceed is to write

$$e^{KS_iS_j} = \cosh(KS_iS_j) + \sinh(KS_iS_j) = \cosh(K) + \sinh(KS_iS_j) = \cosh K + S_iS_j \sinh K, \quad (1)$$

from parity.

- 2) S_2 will later on be a spin to be decimated :

$$\begin{aligned} \sum_{\{S_2\}} e^{KS_1S_2 + KS_2S_3} &= (\cosh K)^2 \sum_{\{S_2\}} (1 + S_1S_2 \tanh K)(1 + S_2S_3 \tanh K) \\ &= 2(\cosh K)^2 [1 + S_1S_3 (\tanh K)^2]. \end{aligned} \quad (2)$$

The terms in S_2^0 and S_2^2 only do survive upon integrating out S_2 .

- 3) The partition function can be written

$$Z(K, N, a) = \sum_{\{S_i\}} \prod_i \exp(KS_iS_{i+1}) = (\cosh K)^N \sum_{\{S_i\}} \prod_i (1 + S_iS_{i+1} \tanh K). \quad (3)$$

Using repeatedly relation (2), for all spins marked with a cross, we arrive at

$$Z(K, N, a) = (\cosh K)^N Z(K', N/b, ba) \quad (4)$$

where since there are $b - 1$ spins integrated out between successive retained spins, it appears that the renormalized model has lattice constant K' such that

$$\boxed{\tanh K' = (\tanh K)^b}. \quad (5)$$

One may note that for $b = 2$, $\tanh K' = (\tanh K)^2 \iff K' = \frac{1}{2} \log \cosh(2K)$, a form that we already met (see the tutorials).

B- The two-dimensional model / take 1

- 4) When computing the partition function, integrating out spins marked with a square in Fig. C1 couples the 4 neighboring spins. Couplings proliferate upon decimation, which is not sustainable.
- 5) Moving the bonds makes the problem locally unidimensional, so that we can make use of the result shown in section A. Therefore, the recursion relation is

$$\boxed{\tanh K' = [\tanh(2K)]^2}. \quad (6)$$

In terms of $x = \tanh K$, this means

$$x' = [\tanh(2K)]^2 = \left[\frac{2 \tanh K}{1 + \tanh^2 K} \right]^2 = \left[\frac{2x}{1 + x^2} \right]^2. \quad (7)$$




FIGURE C1 – Proliferation of couplings under naive renormalization. In one iteration, integrating out the \square spin, the spins marked with circles become coupled by a 4-body term.




FIGURE C2 – Finding the fixed point of the recursion relation (7). The graph is for the $x \mapsto x'$ mapping, with $x = \tanh K$. The non-trivial fixed point is at $x_c \simeq 0.3$.

- 6) Fixed points for relation (7). There are two trivial fixed points : a large-temperature one for $K = 0$, and a low-temperature one for $K \rightarrow \infty$. With $x = \tanh K$, they correspond respectively to $x = 0$ and $x \rightarrow 1$. It can be seen on Fig. C2 that they are both stable, since the derivative is smaller than unity in their vicinity¹. The figure also displays a third (and non-trivial) fixed point, $x_c \simeq 0.3$. For such a value, the $t \mapsto \tanh t$ graph provided in the main text shows that $\tanh t \simeq t$ is a very fair approximation (remember the next Taylor term in the expansion, which is $t^3/3$). Thus, $K_c \simeq 0.3$. A more precise calculation shows that $K_c \simeq 0.305$. This fixed point is unstable, as it should.
- 7) The mean-field prediction is $K_c^{\text{mf}} = 1/4$, since each site has 4 neighbors on the lattice. As always, mean-field overestimates the critical temperature, and correspondingly underestimates the critical K , since it discards fluctuations that destroy order : $K_c^{\text{mf}} < K_c^{\text{exact}}$. Note that Migdal and Kadanoff do a better job here than mean-field.
- 8) Lars Onsager solved the $d = 2$ Ising model in the 1940s. Rudolf Peierls had previously rigorously shown the existence of a phase transition for the $d = 2$ Ising model, in the 1930s.
- 9) Since the large scale features are preserved by renormalization, $\xi' = \xi$, meaning that $\tilde{\xi}'/\tilde{\xi} = 1/b$.
- 10) We know that when $K \rightarrow K'$, $\tilde{\xi} \rightarrow \tilde{\xi}/b$. To loop the loop, we need $\tilde{\xi} \propto |K - K_c|^{-\nu}$. Denoting

1. We are supposed to study the $K \mapsto K'$ mapping, rather than $x \mapsto x'$. Both are equivalent, and if a fixed point is (un)stable in one variable, then so is it for the other. Indeed, let us call f the function behind the mapping $K \rightarrow K' : K' = f(K)$, and we are interested in some K^* with $K^* = f(K^*)$. We change variable to $x = \varphi(K)$, and we take φ to be a bijection (here, a \tanh). Then, $x' = \varphi(K') = \varphi(f(K)) = \varphi(f(\varphi^{-1}(x)))$. Elementary calculus shows that

$$\frac{dx'}{dx} = \varphi'(f(\varphi^{-1}(x))) \frac{f'(\varphi^{-1}(x))}{\varphi'(\varphi^{-1}(x))} = \varphi'(f(K)) \frac{f'(K)}{\varphi'(K)} \implies \left. \frac{dx'}{dx} \right|_{x^*} = \varphi'(f(K^*)) \frac{f'(K^*)}{\varphi'(K^*)} = f'(K^*).$$

This proves that a fixed point exhibits the same stability features in both representations, K or x .

$K = K_c + \delta K$, $K' = K_c + \delta K'$, this means

$$\frac{(\delta K')^{-\nu}}{(\delta K)^{-\nu}} = \frac{\tilde{\xi}'}{\tilde{\xi}} = \frac{1}{b} \implies \boxed{\left. \frac{\partial K'}{\partial K} \right|_{K_c} = b^{1/\nu}}. \quad (8)$$

Differentiating (6), we get

$$(1 - \underbrace{\tanh^2 K'}_{\tanh^4(2K)}) \frac{\partial K'}{\partial K} = 4 \tanh(2K)(1 - \tanh^2 2K) \implies \left. \frac{\partial K'}{\partial K} \right|_{K_c} = \frac{4 \tanh 2K_c}{1 + \tanh^2 2K_c} = 2 \tanh(4K_c), \quad (9)$$

using one more time $\tanh(2t) = \frac{2 \tanh t}{1 + \tanh^2 t}$. Since $K_c \simeq 0.3$, we can read for the graph given in the main text that $\tanh(1.2) \simeq 0.84$. We need to find ν satisfying $2^{1/\nu} \simeq 2 \times 0.84 \simeq 1.68 \simeq 5/3$:

$$\nu \simeq \frac{\log 2}{\log 5/3} \simeq \frac{0.69}{0.51} \simeq \frac{0.69}{0.5(1 + 2 \cdot 10^{-2})} \simeq \frac{0.69}{0.5} (1 - 2 \cdot 10^{-2}) \simeq 1.38 - 2 * 0.14 \simeq 1.35. \quad (10)$$

To conclude, we have found

$$\boxed{\nu \simeq 1.35 \quad \text{while} \quad \nu^{\text{mf}} = \frac{1}{2} \quad \text{and} \quad \nu^{\text{exact}} = 1}. \quad (11)$$

On this count as well, we improve over mean-field.

C- The two-dimensional model / take 2




FIGURE C3 – The 4 steps involved in the procedure, with translation in terms of coupling strengths.

11) We decimate $b - 1$ bonds. According to the result of section A, this means that

$$\boxed{\tanh \tilde{K}_x = (\tanh K_x)^b}, \quad (12)$$

while $\tilde{K}_y = b K_y$.

- 12) The x -bonds are broken and moved as shown in the figure, which then opens the possibility to decimate the y -bonds. Then, the x -bonds undergo a b -fold increase and in the perpendicular direction, $\tanh K'_y = [\tanh(\tilde{K}_y)]^b$. The chain of four moves/decimation is addressed in Fig. C3, where the corresponding change affecting the couplings is sequenced. To summarize :

$$\boxed{K'_x = b \tanh^{-1}(\tanh^b K_x) ; \quad K'_y = \tanh^{-1}(\tanh(bK_y))^b.} \quad (13)$$

- 13) We take $b = 1 + \epsilon$; we will need $\frac{d \tanh^{-1}(x)}{dx} = \frac{1}{1-x^2}$ and $\frac{1}{1-\tanh^2} = \cosh^2$.

$$K'_x = (1 + \epsilon) \tanh^{-1}[(\tanh K_x)^{1+\epsilon}] \quad (14)$$

$$= (1 + \epsilon) \tanh^{-1}[(\tanh K_x)(1 + \epsilon \log \tanh K_x)] + \mathcal{O}(\epsilon^2) \quad (15)$$

$$= (1 + \epsilon) \left\{ \tanh^{-1}(\tanh K_x) + \epsilon \frac{(\tanh K_x) \log \tanh K_x}{1 - \tanh^2 K_x} \right\} + \mathcal{O}(\epsilon^2) \quad (16)$$

$$= K_x + \epsilon \left\{ K_x + \log \tanh K_x \frac{\tanh K_x}{1 - \tanh^2 K_x} \right\} + \mathcal{O}(\epsilon^2). \quad (17)$$

and thus we take, to linear order in ϵ ,

$$K'_x = K_x + \epsilon \{ K_x + (\cosh K_x)(\sinh K_x) \log \tanh K_x \} \quad (18)$$

$$K'_x = \boxed{K_x + \epsilon \left\{ K_x + \frac{\sinh(2K_x)}{2} \log \tanh K_x \right\}}. \quad (19)$$

We treat the recursion relation for K'_y similarly,

$$K'_y = \tanh^{-1} \left\{ \left[\underbrace{\tanh((K_y + \epsilon K_y))}_{\tanh K_y + \epsilon K_y (1 - \tanh^2 K_y)} \right]^{1+\epsilon} \right\} \quad (20)$$

$$= \tanh^{-1} \{ \tanh K_y + \epsilon K_y (1 - \tanh^2 K_y) + \epsilon (\tanh K_y) \log \tanh K_y \} + \mathcal{O}(\epsilon^2) \quad (21)$$

$$= K_y + \epsilon \frac{K_y (1 - \tanh^2 K_y) + (\tanh K_y) \log \tanh K_y}{1 - \tanh^2 K_y}. \quad (22)$$

Finally, we arrive at the same relation as (19)

$$\boxed{K'_y = K_y + \epsilon \left\{ K_y + \frac{\sinh(2K_y)}{2} \log \tanh K_y \right\}}. \quad (23)$$

- 14) With the exact critical coupling $K_c^{\text{exact}} = \frac{1}{2} \log(1 + \sqrt{2})$, one has

$$\exp(2K_c^{\text{exact}}) = 1 + \sqrt{2}, \quad \sinh(2K_c^{\text{exact}}) = 1, \quad \tanh K_c^{\text{exact}} = \frac{1}{1 + \sqrt{2}} \quad (24)$$

which proves the statement : K_c^{exact} is a fixed point of the recursion relation (19).

- 15) We start with Eq. (8), and we denote $F(K)$ the term in the curly brackets in Eq. (19) :

$$(1 + \epsilon)^{1/\nu} = 1 + \frac{\epsilon}{\nu} + \mathcal{O}(\epsilon^2) = \left. \frac{\partial K'}{\partial K} \right|_{K_c^{\text{exact}}} = 1 + \epsilon F'(K_c^{\text{exact}}) \implies \frac{1}{\nu} = F'(K_c^{\text{exact}}). \quad (25)$$

Then,

$$\begin{aligned} F'(K) &= 1 + \cosh(2K) \log \tanh K + \frac{\sinh(2K)}{2} \frac{1 - \tanh^2 K}{\tanh K} = 1 + \cosh(2K) \log \tanh K + 1. \\ \implies F'(K_c^{\text{exact}}) &= 2 - \sqrt{2} \log(1 + \sqrt{2}). \end{aligned} \quad (26)$$

To leading order in ϵ , the critical exponent ν for the correlation length thus reads

$$\boxed{\nu = \frac{1}{2 - \sqrt{2} \log(1 + \sqrt{2})}}. \quad (27)$$

The numerical value of ν , 1.327, is closer to the exact result than its “take 1” counterpart, but not by an impressive amount. Here, the effort did not spectacularly pay off.

D- Generalization to arbitrary dimension (take 1 route)

- 16) On a three-dimensional cubic lattice with $b = 2$, an iteration increases length by a factor of 2, so that $N \rightarrow N/8$. In between retained sites, and before decimation, there are two bonds. To preserve the total “strength” of the lattice, the bond moving step strengthens each bond by a factor 4 (see Fig. C4, where the 4-fold increase is recovered by a different argument). The “take 1” recursion relation between K and K' is thus

$$\boxed{\tanh K' = [\tanh(4K)]^2}. \quad (28)$$




FIGURE C4 – Sketch of the $d = 3$, $b = 2$ procedure. To get the strengthening factor of the bonds, one can focus on “vertical bonds”, and count how many there are, before and after bond moving. Before bond-moving, we have $4 \times 1/4 + 4/2 + 1 = 4$ pairs of bonds, where the weights $1/4$, $1/2$ and 1 come from the fact that these bonds are shared 4 times (along the edges of the cube), 2 times (faces of the cube) or not shared (the central pair of bonds). After bond moving, we have just $4/4 =$ one pair left. This 4-fold decrease in the number of bonds is compensated by a 4-fold increase in the strength. A similar argument applies to bonds in the two other directions (not “vertical”).

- 17) Similarly, for a d -dimensional cubic lattice, N decreases to $N/2^d$ and keeping in mind the factor 2 alluded to in the previous question, we have

$$\boxed{\tanh K' = [\tanh(2^{d-1}K)]^2}. \quad (29)$$

- 18) We start from $K' = \tanh^{-1}[\tanh(2^{d-1}K)]^2$ and use that

$$\tanh^{-1}(x) = \frac{1}{2} \log \left(\frac{1+x}{1-x} \right) \quad \text{together with} \quad 1 - \tanh^2 = \frac{1}{\cosh^2}. \quad (30)$$

In the large K regime, with $x = \tanh(2^{d-1}K) \rightarrow 1$, this yields

$$K' \sim \frac{\log 2}{2} - \frac{1}{2} \log(1 - x^2) \sim \frac{1}{2} \log[\cosh^2(2^{d-1}K)] \sim 2^{d-1} K. \quad (31)$$

Therefore, the small temperature sink is stable for $d > 1$ (and then $K' > K$). This is consistent with the lower critical dimension equal to unity.

- 19) For arbitrary b and d , we start by noting that $N \rightarrow N/b^d$. In between two retained sites, there are b bonds, so that the bond moving step strengthens each bond by a factor b^{d-1} . Decimation then leads to the generalized form

$$\boxed{\tanh K' = [\tanh(b^{d-1}K)]^b}. \quad (32)$$

Migdal-Kadanoff scheme gets worse, when compared to known results, when d increases. One can see in particular that for large d , we get $K_c \sim 2^{2-2d}$, which translates into a d -dependent critical exponent ν for all d . There is no sign of an upper critical dimension. Besides, it can be shown that ν approaches 1 for $d \rightarrow \infty$, which is not the mean-field result.