

SOLUTIONS TD3

The replica method (1/2)

Problem 3.1: CORRELATIONS, p-spin vs REM

① ENERGY CORRELATIONS

We have:
$$\overline{E(\vec{\sigma})E(\vec{\tau})} = \sum_{i_1 < i_2 < \dots < i_p \in N} \sum_{j_1 < j_2 < \dots < j_p \in N} \overline{J_{i_1 \dots i_p} J_{j_1 \dots j_p}} \sigma_{i_1} \dots \sigma_{i_p} \tau_{j_1} \dots \tau_{j_p}$$

The random couplings are independent, and therefore the average is non-zero only whenever all the indices are the same. Using the expression of the Variance:

$$\overline{E(\vec{\sigma})E(\vec{\tau})} = \sum_{i_1 < \dots < i_p \in N} \frac{p!}{N^{p-1}} \sigma_{i_1} \tau_{i_1} \sigma_{i_2} \tau_{i_2} \dots \sigma_{i_p} \tau_{i_p}$$

Now, the constraint on the non-repeating indices can be released using that:

$$\sum_{i_1 < \dots < i_p} \approx \frac{1}{p!} \sum_{i_1, \dots, i_p}$$

and thus $\overline{E(\vec{\sigma})E(\vec{z})} = N \left(\sum_{i_2=1}^N \frac{\sigma_{i_2} z_{i_2}}{N} \right) \dots \left(\sum_{i_p=1}^N \frac{\sigma_{i_p} z_{i_p}}{N} \right) = N [q(\vec{\sigma}, \vec{z})]^p$

where $q = q(\vec{\sigma}, \vec{z})$ is the overlap.

One has $q \leq 1$ (Cauchy-Schwartz): therefore, when $p \rightarrow \infty$ $q^p \rightarrow 0$, and correlation vanish as in the REM.

Problem 3.2: THE ANNEALED FREE-ENERGY

1 ENERGY CONTRIBUTION

The averaged partition function is:

$$\overline{Z} = \int_{S^N} d\vec{\sigma} \overline{e^{-\beta E(\vec{\sigma})}} = \int_{S^N} d\vec{\sigma} \overline{e^{-\beta \sum_{i_2 < \dots < i_p} \sigma_{i_2} \dots \sigma_{i_p}}} \stackrel{\text{Independence}}{=} \int_{S^N} d\vec{\sigma} \prod_{i_2 < \dots < i_p} \overline{e^{-\beta \sum_{i_2 < \dots < i_p} \sigma_{i_2} \dots \sigma_{i_p}}}$$

where $\int_{S^N} d\vec{\sigma}$ is the integral on the surface of the sphere in dimension N .

By definition, $f_a = \lim_{N \rightarrow \infty} \frac{1}{N} \log \overline{Z}$.

Performing the Gaussian integral (e.g. by completing the square) we get:

$$\overline{e^{-\beta \sum_{i=1}^p \sigma_i^2}} = e^{\frac{\beta^2}{2} \sigma_{i_1}^2 \dots \sigma_{i_p}^2} \frac{p!}{N^{p-1}} \quad \text{and thus:}$$

$$\overline{Z} = \int_{S_N} d\vec{\sigma} e^{\frac{\beta^2}{2} N \left(p! \prod_{i_1 < \dots < i_p} \frac{\sigma_{i_1}^2}{N} \dots \frac{\sigma_{i_p}^2}{N} \right)} = e^{\frac{\beta^2 N}{2}} \int_{S_N} d\vec{\sigma}.$$

Using that $\sum_i \sigma_i^2 = N$

2] ENTROPY CONTRIBUTION

The Stirling formula implies $\left(\frac{N}{2}\right)! \stackrel{N \gg 1}{\approx} e^{-\frac{N}{2}} \left(\frac{N}{2}\right)^{N/2} = e^{-\frac{N}{2} + \frac{N}{2} \log\left(\frac{N}{2}\right)}$

and thus

$$\int_{S_N} d\vec{\sigma} = \frac{(\pi N)^{N/2}}{\left(\frac{N}{2}\right)!} \approx e^{\frac{N}{2} [\log(\pi N) + 1 - \log\left(\frac{N}{2}\right)] + o(N)}$$

$$= e^{\frac{N}{2} \log(2\pi e) + o(N)}$$

Putting everything together, we find:

$$\overline{Z} = \exp \left\{ N \left(\frac{\beta^2}{2} + \frac{1}{2} \log(2\pi e) \right) + o(N) \right\}$$

and therefore

$$f_a = -\frac{1}{\beta} \left[\frac{\beta^2}{2} + \frac{1}{2} \log(2\pi e) \right]$$

The difference comes from the entropic contribution $\int_{S^N} d\vec{\sigma}$, and it is due to the fact that the phase space of the spherical model is different from that of the REM, where spins are discrete variables ± 1 .

Problem 3.3: QUENCHED FREE-ENERGY, REPLICAS

① STEP 1: FROM QUENCHED RANDOMNESS TO INTERACTIONS

The n -th power of the partition function is:

$$Z^n = \left(\prod_{a=1}^n \int_{S^N} d\vec{\sigma}^a \right) \exp \left[-\beta \sum_{i_1 < \dots < i_p} J_{i_1 \dots i_p} (\sigma_{i_1}^1 \dots \sigma_{i_p}^1 + \sigma_{i_1}^2 \dots \sigma_{i_p}^2 + \dots + \sigma_{i_1}^n \dots \sigma_{i_p}^n) \right]$$

When averaging over the couplings $J_{i_1 \dots i_p}$, we use again the properties of INDEPENDENCE and GAUSSIANITY and get:

$$\overline{Z^n} = \left(\prod_{a=1}^n \int_{S^N} d\vec{\sigma}^a \right) \prod_{i_1 < \dots < i_p} e^{-\frac{\beta^2}{2} \frac{p!}{N^{p-1}} (\sigma_{i_1}^1 \dots \sigma_{i_p}^1 + \sigma_{i_1}^2 \dots \sigma_{i_p}^2 + \dots + \sigma_{i_1}^n \dots \sigma_{i_p}^n)^2}$$

The square at the exponent can be re-written as:

$$\sum_{a=1}^n \sum_{b=1}^n (\sigma_{i_2}^a \sigma_{i_2}^b) \dots (\sigma_{i_p}^a \sigma_{i_p}^b)$$

Therefore, using again that $\sum_{i_2 < \dots < i_p} \simeq \frac{1}{p!} \sum_{i_2, \dots, i_p}$

we obtain:

$$\begin{aligned} \overline{Z^n} &= \left(\prod_{a=1}^n \int_{S^N} d\vec{\sigma}^a \right) e^{\frac{\beta^2}{2} N \sum_{a,b=1}^n \sum_{i_2, i_2, \dots, i_p} \frac{(\sigma_{i_2}^a \sigma_{i_2}^b)}{N} \dots \frac{(\sigma_{i_p}^a \sigma_{i_p}^b)}{N}} \\ &= \left(\prod_{a=1}^n \int_{S^N} d\vec{\sigma}^a \right) e^{\frac{\beta^2}{2} N \sum_{a,b=1}^n \left(\frac{\vec{\sigma}^a \cdot \vec{\sigma}^b}{N} \right)^p} \quad (*) \end{aligned}$$

In this expression, the quenched randomness has disappeared, but the replicas are coupled!

Step 1: start from expression with replicas decoupled, subject to same disorder.
After averaging, end up with COUPLED REPLICAS (interacting theory), no disorder.

2 STEP 2: EMERGING ORDER PARAMETERS

The final expression of $\overline{z^n}$ shows that the integrand depends on the variables σ_i^a only through global quantities, the scalar products between the $\vec{\sigma}^a$.

We can therefore identify a set of functions, the overlaps between the replicas:

$$q^{ab} = q(\vec{\sigma}^a, \vec{\sigma}^b) = \frac{\sum_{i=1}^N \sigma_i^a \sigma_i^b}{N},$$

that are ORDER PARAMETERS of the theory, like the magnetization $m = \frac{1}{N} \sum_{i=1}^N \sigma_i$ in the mean-field Ising model. In particular, in (*) we can replace the integral over all possible configurations of the $\vec{\sigma}^a$ with an integral over all possible values of the overlaps, using:

$$\int \prod_{a < b} dq_{ab} \delta(q(\vec{\sigma}^a, \vec{\sigma}^b) - q_{ab}) = 1$$

↑
integration variables, numbers

↖
functions

Plugging this in (*) we obtain:

$$\left(\prod_{a=1}^n \int_{S_N} d\vec{\sigma}^a \right) e^{\frac{\beta^2}{2} N \sum_{a,b=1}^n \left(\frac{\vec{\sigma}^a \cdot \vec{\sigma}^b}{N} \right)^p}$$

$$= \left(\prod_{a=1}^n \int_{S_N} d\vec{\sigma}^a \right) \left(\int \prod_{a < b} dq_{ab} \delta(q(\vec{\sigma}^a, \vec{\sigma}^b) - q_{ab}) \right) e^{\frac{\beta^2}{2} N \sum_{a,b=1}^n \left(\frac{\vec{\sigma}^a \cdot \vec{\sigma}^b}{N} \right)^p}$$

↔ exchange

$$= \int \prod_{a < b} dq_{ab} \left[\left(\prod_{a=1}^n \int_{S_N} d\vec{\sigma}^a \right) \prod_{a < b} \delta(q(\vec{\sigma}^a, \vec{\sigma}^b) - q_{ab}) \right] e^{\frac{\beta^2}{2} N \sum_{a,b=1}^n q_{ab}^p}$$

We call

$$V(\{q_{ab}\}_{a < b}) = \left(\prod_{a=1}^n \int_{S_N} d\vec{\sigma}^a \right) \prod_{a < b} \delta(q(\vec{\sigma}^a, \vec{\sigma}^b) - q_{ab}) = e^{NS[\{q_{ab}\}] + d(N)}$$

where $S[\cdot]$ is the entropy of configurations satisfying the constraint on the overlaps being equal to q_{ab} .

We introduce the $n \times n$ matrix with components:

$$Q_{ab} = \begin{cases} q_{ab} & a < b \\ 1 & a = b \\ q_{ba} & b < a \end{cases} \quad \begin{array}{l} \text{OVERLAP} \\ \text{MATRIX} \end{array}$$

Then it can be shown (exercise! solution below) that

$$V[Q] = e^{N S[Q] + o(N)}, \quad S[Q] = \frac{n}{2} \log(2\pi e) + \frac{1}{2} \log \det[Q]$$

and thus:

$$\bar{Z}^n = \int \prod_{a < b} dq_{ab} e^{N \left\{ \frac{\beta^2}{2} \sum_{a,b} q_{ab}^P + \frac{n}{2} \log(2\pi e) + \frac{1}{2} \log \det[Q] \right\}}$$

This theory now is expressed only in terms of Q :

$$\bar{Z}^n = \int \prod_{a < b} dq_{ab} e^{N A_n[Q] + o(N)}$$

"the action"

$$A_n[Q] = \frac{\beta^2}{2} \sum_{a,b} q_{ab}^P + \frac{n}{2} \log(2\pi e) + \frac{1}{2} \log \det[Q]$$

Step 2: re-write the integral over configurations as integral over emerging order parameter q_{ab} . Same as magnetization for mean-field Ising:

$$Z_{\text{Ising}} = \sum_{\vec{s}=1}^{2^N} e^{-\frac{\beta J}{N} \sum_{i,m} s_i s_m} \longrightarrow \sum_M \underbrace{N(m)}_{\substack{\text{number of configurations} \\ \text{with magnetization } m}} e^{-\beta J N m^2}$$

$\langle s_i \rangle = \frac{1}{N} \sum_i s_i$

In the replica calculation, have $\frac{n(n-1)}{2}$ order parameters to integrate over. We started with Nn variables: HUGE DIMENSIONALITY REDUCTION due to mean-field.

[3] STEP 3: SADDLE-POINT, SELECTING THE TYPICAL

For large N , the integral over the space of $n \times n$ matrices Q can be computed with a saddle-point approximation.

The derivative with respect to a matrix Q has to be intended as the derivative wrt. its components:

$$\frac{\partial \sum_{c,d} q_{cd}^p}{\partial q_{ab}} = 2p q_{ab}^{p-1} \quad , \quad \frac{\partial \log \det[Q]}{\partial q_{ab}} = 2(Q^{-1})_{ab}$$

↑ symmetry Q

Therefore the saddle point equations read

$$\beta^2 p q_{ab}^{p-1} + (Q^{-2})_{ab} \Big|_{Q=Q^*} = 0 \quad \text{for all } a < b \quad (**)$$

Q^* = saddle-point value

To proceed, need to make assumptions on structure of q_{ab} at the saddle point: a "VARIATIONAL ANSATZ".

Comment: there is one simple solution to the saddle point equations:

$$Q_a^* = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} = \mathbb{1}^{n \times n}$$

This solves trivially the equations since both terms in (**) are zero.

In this case:

$$\overline{Z}^n = e^{NA_n[Q_n^*] + o(N)} = e^{Nn \left\{ \frac{\log(2\pi e)}{2} + \beta^2/2 \right\} + o(N)}$$

Using the replica trick:

$$f = -\frac{1}{\beta} \lim_{N \rightarrow \infty} \lim_{n \rightarrow 0} \frac{\overline{Z}^n - 1}{Nn} = -\frac{1}{\beta} \left\{ \frac{\log(2\pi e)}{2} + \beta^2/2 \right\}$$

Same result as the annealed.

At high T , this is the good solution.

There is a critical temperature T_c there is another solution, with lower free energy than the above!

Extra: Volume Term

$$\begin{aligned}
 V &= \left(\prod_{a=1}^n \int d\vec{\sigma}^a \right) \prod_{a < b} \delta(q(\vec{\sigma}^a, \vec{\sigma}^b) - q_{ab}) \prod_a \delta(q(\vec{\sigma}^a, \vec{\sigma}^a) - 1) \\
 &= N^{\frac{n(n-1)}{2} + n} \left(\prod_{a=1}^n \int d\vec{\sigma}^a \right) \prod_{a < b} \delta(\vec{\sigma}^a \cdot \vec{\sigma}^b - N q_{ab}) \prod_a \delta(\vec{\sigma}^a \cdot \vec{\sigma}^a - N) \\
 &= N^{\frac{n(n+1)}{2}} \left(\prod_{a=1}^n \int d\vec{\sigma}^a \right) \int \prod_{a \leq b} \frac{d\lambda_{ab}}{\sqrt{2\pi}} e^{i \sum_{a \leq b} \lambda_{ab} (\vec{\sigma}^a \cdot \vec{\sigma}^b - q_{ab} N)}
 \end{aligned}$$

Where $Q_{ab} = \begin{cases} q_{ab} & \text{if } a < b \\ 1 & \text{if } a = b \\ q_{ab} & \text{if } b < a \end{cases}$

$$= \left(\frac{N}{\sqrt{2\pi}} \right)^{\frac{n(n+1)}{2}} \int \prod_{a \leq b} d\lambda_{ab} e^{-iN \sum_{a \leq b} \lambda_{ab} Q_{ab}} \left[\int \prod_{a=1}^n d\vec{\sigma}^a e^{i \sum_{a \leq b} \lambda_{ab} \vec{\sigma}^a \cdot \vec{\sigma}^b} \right]$$

Now, $\sum_{a \leq b} = \frac{1}{2} \sum_{a \neq b} (\dots) + \sum_a (\dots)$

Call $\tilde{\lambda}_{ab} = \begin{cases} -\frac{i \lambda_{ab}}{2} & a \neq b \\ -i \lambda_{aa} & a = b \end{cases}$

And get:

$$= \left(\frac{N}{\sqrt{2\pi}} \right)^{\frac{n(n+1)}{2}} \int \prod_{a \leq b} d\lambda_{ab} e^{N \sum_{a,b} \tilde{\lambda}_{ab} Q_{ab}} \left[\int \prod_{a=1}^n d\vec{\sigma}^a e^{-\frac{1}{2} \sum_{a,b} 2\tilde{\lambda}_{ab} \vec{\sigma}^a \cdot \vec{\sigma}^b} \right]$$

Now, the integral over the variables $\vec{\sigma}_i$ is a multivariate Gaussian integral. One has:

$$\begin{aligned} I &= \int \prod_{a=1}^n \prod_{i=1}^N d\sigma_i^a e^{-\frac{1}{2} \sum_{a,b} \sum_{ij} \sigma_i^a (2\tilde{\lambda}_{ab} \delta_{ij}) \sigma_j^b} \\ &= (2\pi)^{\frac{Nn}{2}} \left[\det(2\tilde{\Lambda}) \right]^{-\frac{N}{2}} = e^{\frac{Nn}{2} \log(2\pi) - \frac{N}{2} \log(\det[2\tilde{\Lambda}])} \end{aligned}$$

Now, we are left with:

$$\begin{aligned} V &= \left(\frac{N}{\sqrt{2\pi}} \right)^{\frac{n(n+1)}{2}} (2\pi)^{\frac{Nn}{2}} \int \prod_{a \leq b} d\lambda_{ab} e^{N \sum_{a,b} \tilde{\lambda}_{ab} Q_{ab} - \frac{N}{2} \log \det[2\tilde{\Lambda}]} \\ &= C_{N,n} (2\pi)^{\frac{Nn}{2}} \int \prod_{a \leq b} d\tilde{\lambda}_{ab} e^{N \text{tr}(\tilde{\Lambda} Q) - \frac{N}{2} \log \det[2\tilde{\Lambda}]} \end{aligned}$$

This is an integral in matrix space that can be performed with a saddle point, which gives:

$$Q - \frac{1}{2}(\tilde{\Lambda})^{-1} = 0 \Rightarrow \tilde{\Lambda}^* = (2Q)^{-1}$$

$$\det [2\tilde{\Lambda}] \Big|_{\tilde{\Lambda}=\tilde{\Lambda}^*} \Rightarrow \det [Q^{-1}] = (\det Q)^{-1}$$

Putting everything together:

$$V = e^{\frac{Nn}{2} \log(z\pi) + \frac{Nn}{2} + \frac{N}{2} \log \det(Q)}$$

$$= e^{\frac{Nn}{2} \log(2\pi e) + \frac{N}{2} \log \det(Q)}$$

One can show that with this choice, the replica calculation reproduces the annealed calculation we did in Problem 1 \rightarrow EXERCISE

However, this is wrong in the low- T phase! There, fluctuations dominate and have to be captured by another structure of the matrix: 1RSB.

Assumption 2:
1-step RSB

$$(q_{ab})_{ab} = \left(\begin{array}{c} \overleftarrow{m} \\ \begin{array}{|c|} \hline 1 \quad q \\ \hline q \quad 1 \\ \hline \end{array} \\ \begin{array}{|c|} \hline 1 \quad q \\ \hline q \quad 1 \\ \hline \end{array} \\ \begin{array}{|c|} \hline 1 \quad q \\ \hline q \quad 1 \\ \hline \end{array} \\ \hline \end{array} \right)$$

meaning: assume that n replicas organize in n/m groups of size m , with them falling in similar configurations at overlap q .

Notice: m is arbitrary parameter, to be optimized-over in saddle point.