

PROBLEM 8

Anderson on Bethe Lattice (1/2)

Problem 8 : BETHE LATTICE, RECURSIONS, CAVITY

GREEN FUNCTIONS IDENTITIES

By definition,

$$G = (z - H)^{-1} = [(z - H_0)(1 - (z - H_0)^{-1} H_1)]^{-1} = (1 - (z - H_0)^{-1} H_1)^{-1} G^0$$


Multiplying to the left by $1 - (z - H_0)^{-1} H_1$ we get:

$$(1 - G^0 H_1) G = G^0 \Rightarrow G - G^0 H_1 G = G^0 \Rightarrow G = G^0 H_1 G + G^0$$

When iterated, this relation gives rise to the perturbative series for G :

$$G = G^0 + G^0 H_1 G^0 + G^0 H_1 G^0 H_1 G^0 + \dots$$

2] CAVITY EQUATIONS



In this case, the term H_1 corresponds to the three links in pink in the figure.

Removing those links, one is decoupling the root from the $(K+1)$ subtrees with vertex a_1, \dots, a_K .

In particular, $H_0 = W \cdot V_0 |0\rangle\langle 0| + \sum_{i=1}^K H_i^{(0)}$ where $H_i^{(0)}$ is the Hamiltonian restricted to the subtree with vertex a_i , $i=1, \dots, K+1$. Since each subtree is completely disconnected with the root, the Green function G_{ai}^{cav} depends only on the Hamiltonian restricted to the subtree: it is the same that one would get removing the site 0.

The Green function relation can be iterated at any order.
Let us go to 2nd order in H_1

$$G = G_0 + G_0 H_1 G_0 + G_0 H_1 G_0 H_1 G$$

Let us take matrix elements:

$$G_{00} = G_{00}^0 + \sum_{a,b} G_{0a}^0 (H_1)_{ab} G_{b0}^0 + \sum_{a,b,c,d} G_{0a}^0 (H_1)_{ab} G_{bc}^0 (H_1)_{cd} G_{d0}^0$$


Now, G_{0a}^0 is non zero only for $a=0$, and $(H_1)_{00}=0$: the first order vanishes. Also, $(H_1)_{0a}$ is non-zero only if $a \in \{a_1, \dots, a_{k+1}\}$, the neighbors of 0, and it equals to $-t_{0a}$.

Thus:

$$G_{00} = G_{00}^0 + \sum_{i=1}^{k+1} G_{00}^0 t_{0a_i} G_{a_i a_i}^0 t_{a_i 0} G_{00} \Rightarrow G_{00} = \left(1 - G_{00}^0 \sum_{i=1}^{k+1} t_{0a_i}^2 G_{a_i a_i}^{cav} \right)^{-1} G_{00}^0$$

Using that $G_{00}^0 = (z - W V_0)^{-1}$, we get the first equation.

Let us iterate this procedure: we consider a subtree with origin in a_i , and define H_1 the links connecting the origin to the "descendents":




Repeating the steps above, we find:

$$G_{a_i}^{cav} = \frac{1}{z - W V_{a_i} - \sum_{b \in \partial a_i} t_{a_i b}^2 G_b^{cav}} \quad \begin{array}{c} \text{by definition of } \sigma^{cav} \\ \downarrow \\ \equiv \end{array} \frac{1}{z - W V_{a_i} - \sigma_{a_i}^{cav}}$$

this sum is now over K sites, not $K+1$

$$\text{Then } \sigma_{a_i}^{cav} = \sum_{b \in \partial a_i} t_{a_i b}^2 G_b^{cav} \quad \begin{array}{c} \text{use def of } \sigma^{cav} \\ \downarrow \end{array} = \sum_{b \in \partial a_i} t_{a_i b}^2 \frac{1}{z - W V_b - \sigma_b^{cav}}$$

[3] EQUATIONS FOR THE DISTRIBUTION



The functions $G_{b_1}^{\text{cav}}$ and $\sigma_{b_1}^{\text{cav}}$ depend only on the sites (and on the randomness V_i) in the subtree T_1 , which is not overlapping with the subtree T_2 .

Therefore, the two random function are independent.

Moreover, they are statistically equivalent (the sub-trees are statistically identical), and so they can be considered as identically distributed variables.

[4] THE "LOCALIZED" SOLUTION

- We have $\sigma_a^{\text{cav}} = R_a - i\Gamma_a$

The cavity equation becomes:

$$R_{a_i}(E+i\eta) - i\Gamma_{a_i}(E+i\eta) = \sum_{b \in \partial a_i} t_{a_i b}^2 \frac{1}{[E - W V_{a_i} - R_b(E+i\eta)] + i[\eta + \Gamma_b(E+i\eta)]}$$

$$\downarrow$$

$$\sum_{b \in \partial a_i} t_{a_i b}^2 \frac{E - W V_{a_i} - R_b(E+i\eta) - i[\eta + \Gamma_b(E+i\eta)]}{[E - W V_{a_i} - R_b(E+i\eta)]^2 + [\eta + \Gamma_b(E+i\eta)]^2}$$

Equating real and imaginary part, we get the two equations.

- The equation for P_a is satisfied, for $\eta=0$, setting $P_a = P_b = 0$.

The solution $P=0$ corresponds to localization. It is always a solution when $\eta=0$.

- The Anderson criterion states that in the localized phase, when $\eta \downarrow 0$ the distribution of P tends to $\delta(P)$.

This means that the solution $P=0$, which holds for $\eta=0$, remains a stable solution also when adding $\eta>0$, taking $N \rightarrow \infty$ and then switching off η .

TO ESTABLISH LOCALIZATION, WE
HAVE TO STUDY THE STABILITY OF
THE SOLUTION $P(P) = \delta(P)$.